

32802-67

4/10/2015

1/18



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

James McFadden, Ph.D.
Director, Regulatory Affairs
Howard Johnson's Enterprises, Inc.
9675 S. 60th Street
Franklin, WI 53132

APR 10 2015

Subject: Label Notification per PRN 98-10 – Adding Required Language for Florida
Crabgrass Preventer Plus with .58 Barricade Preemergence Herbicide
EPA Registration Number: 32802-67
Application Date: February 13, 2015
Decision Number: 500593

Dear Dr. McFadden:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped "Notification" and will be placed in our records. If you have any questions, you may contact Terri Stowe at (703) 305-6117 or via email at stowe.terri@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Kathryn V. Montague".

Kathryn V. Montague
Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

2/18

CRABGRASS PREVENTER PLUS WITH .58% BARRICADE® PREEMERGENCE HERBICIDE

Sublabel A: Residential lawns and outdoor domestic use with applications by homeowners]

FOR SELECTIVE PREEMERGENCE CONTROL OF GRASS and BROADLEAF WEEDS IN RESIDENTIAL LAWNS

STOPS MANY [UGLY] WEED GRASSES BEFORE THEY GROW. PREVENTS CRABGRASS, ANNUAL BLUEGRASS, FOXTAIL & MORE.

(See Marketing Claims Section for Additional Claims)

ACTIVE INGREDIENT:
Prodiamine: (CAS 29091-21-2) 0.58%
OTHER INGREDIENTS: 99.42%
TOTAL: 100.00%

NOTIFICATION

APR 10 2015

EPA Reg. No. 32802-67
EPA Est. No. 59144-GA-1 (A), 32802-MO-1 (B),
37351-CO-1 (C), 8378-IN-1 (D), 9404-FL-1 (E)
First letter of bag code shows establishment where manufactured..
[See tag attached to bag for fertilizer guaranteed analysis.]

GUARANTEED ANALYSIS XX-X-X	
Total Nitrogen (N)	XX%
x.x% ammoniacal nitrogen	
x.x% urea nitrogen*	
Available Phosphate (P ₂ O ₅)	X%
Soluble Potash (K ₂ O)	X%
Sulfur (S)	X%
x.x% free sulfur (S)	
Derived from: polymer-coated sulfur-coated urea, urea, ammonium phosphate, potassium chloride. *A portion of the urea has been coated to provide x.x% coated slow-release nitrogen. This product and/or its manufacture is covered by one or more of the following U.S. patents and international counterparts thereof: US Patent Numbers: xxxxxx	
Note: Fertilizer grades with nutrient contents expressed as the percentages of N, P ₂ O ₅ and K ₂ O may include the following: 23-3-7 or x-x-x. The fertilizer grade analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.	

4/18

First Aid	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or physician immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or physician. • Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
Hot line number	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.	

USE PRECAUTIONS: The following precautions apply to the use of this product in residential lawns. Application of this product may thin emerged annual bluegrass and newly overseeded grasses. Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil). Do not apply to newly set sod until the following year. Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural practices (such as core aeration) may result in reduced weed control. Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn and sweep any product that lands on the driveway, sidewalk or street back onto your lawn.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in established residential lawns and landscape ornamentals and in established perennials and wildflower plantings in residential settings. It controls susceptible weeds by inhibiting weed seed germination and root development. When used according to directions, it will control: barnyardgrass; bluegrass, annual (*Poa annua*); carpetweed; chickweed, common; chickweed, mouseear (from seed); crabgrass (large, smooth); crowfootgrass; cupgrass, woolly; foxtails, annual; goosegrass¹; henbit; itchgrass; johnsongrass (from seed); junglerice; knotweed; kochia; lambsquarter, common; lovegrass; panicum (Texas, fall, browntop); pigweed; purslane, common; pusley, Florida; rescuegrass²; shepherd's purse²; signalgrass, broadleaf; speedwell, Persian; sprangletop; spurge, prostrate; witchgrass; woodsorrel, yellow (from seed).

c c c c c c
c c c c c c
c c c c c c
c c c c c c
c c c c c c
c c c c c c
c c c c c c
c c c c c c

1. In many areas [where not prohibited by fertilizer use restrictions**] a single application of 2.5 to 6 pounds / 1000 square feet (112 to 259 pounds/acre) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 2.5 to 3.9 pounds / 1000 square feet (112 to 173 pounds/acre) [to areas not prohibited by fertilizer use restrictions,] followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see "Minimum and Maximum Application Rate by Turfgrass Species" table in Rates of Application section of this label).

2 For this weed, apply in late summer, fall or winter prior to germination.

3 Suppression only.

[**Consult your State Cooperative Extension Service for current information on annual application limitations for lawn fertilizers in your region.]

5/18


WHEN TO APPLY TO ESTABLISHED TURF: This product must be applied prior to germination of weed seed, as it will not control established weeds.

[Graphic – Map showing approximate crabgrass germination dates]

APPLICATION DIRECTIONS: Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days after application.

[BROADCAST SPREADERS: For best results, apply in a circular path starting on the outside and working toward the center overlapping slightly to insure complete coverage.

DROP SPREADERS: Apply 2 strips along each end. Then apply along length of lawn, overlapping slightly and shutting off the spreader before each turn at edge of border strips.



SPREADER SETTINGS to apply at the listed pounds per 1000 square feet					
	[1.9 lbs]	[2.5 lbs]	[3.16 lbs]	[4 lbs]	[6 lbs]
Spreader A					
<i>(May contain additional spreader brands/models, spreader settings and spreader use instructions as appropriate for the product)</i>					

SETTING YOUR LAWN SPREADER

THESE SETTINGS ARE A GUIDE. They were determined by landscape professionals using new spreaders. The condition of your spreader, your walking speed, the roughness of the lawn, and even the humidity can affect how fast the lawn food comes out. Watch flow rate. If fertilizer appears to be coming out either slower or faster than you are covering your lawn, adjust the setting. Close the spreader when filling, turning or stopping.

[Advisory For Fertilizers Containing Iron:
 Keep this product off sidewalks, patios, driveways, etc. This fertilizer may stain concrete and painted surfaces. If it comes in contact with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.]

(Optional/Alternate Application Directions)

1. Determine size of area to be treated - measure length and width in feet, then multiply to determine coverage area in square feet (length x width). Next, subtract non-lawn areas.
2. Use the spreader setting chart to determine the appropriate setting.
3. Apply evenly over dry grass, avoiding streaks, skips and overlaps - thorough coverage is important. On hot days (above 80° F), water the lawn to rinse particles off the grass.

Note: The spreader settings listed at right are approximate. It is important to check calibration of spreader frequently to ensure spreader is functioning properly and applying product evenly. To check calibration, accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust spreader setting if necessary in order to ensure correct rate of application.

9/18

Manufactured by: Howard Johnson's Enterprises, Inc.
9675 S 60th Street, Franklin, WI 53132

BARRICADE" is a registered trademark of Syngenta Crop Protection

Read the entire "Directions for Use" and "Warranty Statements" before using this product. If terms are not acceptable, return the unopened product container at once to the seller.

(Brackets [] contain optional language for use in states where the maximum application rate is limited by N restrictions)

NET WEIGHT: XXX LB

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See back panel for further Precautions and First Aid information.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.

First Aid	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or physician immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or physician. • Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
Hot line number	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.	

ENVIRONMENTAL HAZARDS:

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

10/18

USE PRECAUTIONS: The following precautions apply to the use of this product in turf grasses and lawns. Application of this product may thin emerged annual bluegrass and newly overseeded grasses. Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil). Do not apply to newly set sod until the following year. Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural practices (such as core aeration) may result in reduced weed control. Do not apply to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species. Do not apply through any type of irrigation system. Do not apply aerially. Do not graze or feed livestock forage cut from treated areas. Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn and sweep any product that lands on the driveway, sidewalk or street back onto your lawn.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

(Alternate wording for WPS exclusionary statement, for use when the product labeling contains only turf uses, (ie when label is split to exclude the directions for use on ornamental sites.))

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in established lawns and ornamental turf (excluding golf course putting greens), landscape ornamentals and established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seed germination and root development. When used according to directions, it will control: barnyardgrass; bluegrass, annual (*Poa annua*); carpetweed; chickweed, common; chickweed, mouseear (from seed); crabgrass (large, smooth); crowfootgrass; cupgrass, woolly; foxtails, annual; goosegrass¹; henbit; itchgrass; johnsongrass (from seed); junglerice; knotweed; kochia; lambsquarter, common; lovegrass; panicum (Texas, fall, browntop); pigweed; purslane, common; pusley, Florida; rescuegrass³; shepherd's purse²; signalgrass, broadleaf; speedwell, Persian; sprangletop; spurge, prostrate; witchgrass; woodsorrel, yellow (from seed).

1. In many areas [where not prohibited by fertilizer use restrictions**] a single application of 2.5 to 6 pounds / 1000 square feet (112 to 259 pounds/acre) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 2.5 to 3.9 pounds / 1000 square feet (112 to 173 pounds/acre) [to areas not prohibited by fertilizer use restrictions,] followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see "Minimum and Maximum Application Rate by Turfgrass Species" table in Rates of Application section of this label).

2 For this weed, apply in late summer, fall or winter prior to germination.

3 Suppression only.

[**Consult your State Cooperative Extension Service for current information on annual application limitations for lawn fertilizers in your region.]

WHEN TO APPLY TO ESTABLISHED TURF: This product must be applied prior to germination of weed seed, as it will not control established weeds.

[Graphic – Map showing approximate crabgrass germination dates]

APPLICATION DIRECTIONS: Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days after application.

RATES OF APPLICATION: This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not exceed the maximum yearly rate listed:

Minimum and Maximum Application Rates (by Turf Grass Species) ¹		
Turf Species	Minimum Rate	Maximum Rate
Bahiagrass ^[2] , Centipedegrass ^[2] Bermudagrass ³ , Zoysiagrass Seashore Paspalum St. Augustinegrass Tall-Fescue (including turf-type)	4 pounds per 1000 sq. ft (1.75 lbs/acre)	6 pounds/1000 sq. ft. per year (259 lbs/acre per year)
Buffalograss Kentucky Bluegrass Perennial Ryegrass	2 pounds per 1000 sq. ft (87.5 lbs/acre)	4 pounds/1000 sq. ft. per year (173 lbs/acre per year)
Creeping Red Fescue	1.5 pounds per 1000 sq. ft (65.3 lbs/acre)	2.9 pounds per 1000 sq. ft. per year (260 lbs/acre per year)
Creeping Bentgrass	1.5 pounds per 1000 sq. ft (65.3 lbs/acre)	2.5 pounds per 1000 sq. ft. per year (112 lbs/acre per year)
<p>1 These are the maximum rates per calendar year.</p> <p>2 [In Florida: Do not exceed 2 applications per year to Bahiagrass and Centipedegrass in the north. Do not exceed 3 applications per year to Centipedegrass in the central and south regions. Consult your State Cooperative Extension Service for current information on annual application limitations for lawn fertilizers in your region.]</p> <p>3 May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 2 lbs / 1000 square feet (87.5 pounds/acre). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.</p>		

(Optional/Alternate Application Directions)

1. Determine size of area to be treated - measure length and width in feet, then multiply to determine coverage area in square feet (length x width). Next, subtract non-lawn areas.
2. Use the spreader setting chart to determine the appropriate setting.
3. Apply evenly over dry grass, avoiding streaks, skips and overlaps - thorough coverage is important. On hot days (above 80° F), water the lawn to rinse particles off the grass.

Note: The spreader settings listed [below][at right] are approximate. It is important to check calibration of spreader frequently to ensure spreader is functioning properly and applying product evenly. To check calibration, accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust spreader setting if necessary in order to ensure correct rate of application.

SPREADER SETTINGS to apply at the listed pounds per 1000 square feet					
	[1.9 lbs]	[2.5 lbs]	[3.16 lbs]	[4 lbs]	[6 lbs]
Spreader A					
(May contain additional spreader brands/models, spreader settings and spreader use instructions as appropriate for the product)					

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control.

WHEN TO OVERSEED AFTER APPLICATION: This product will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/ reseeding results.

Use Rate	Months Before Overseeding		
	North	Transition	South
Lbs/1000 sq ft			
1.9 lbs/1000 sq ft (86 pounds/acre)	4	4	4
2.5 lbs/1000 sq ft (112 pounds/acre)	5	4	4
2.9 lbs/1000 sq ft (130 pounds/acre)	6	5	5
3.16 lbs/1000 sq ft (138 pounds/acre)		6	6
4 lbs/1000 sq ft (173 pounds/acre)		7	7
4.5 lbs/1000 sq ft (197 pounds/acre)			9
5.14 lbs/1000 sq ft (224 pounds/acre)			10
5.94 lbs/1000 sq ft (259 pounds/acre)			12

LANDSCAPE ORNAMENTALS (including established perennials and wildflower plantings): This product may be applied for residual preemergence weed control in ornamentals. Apply at the rate of 2.5 to 5.94 pounds / 1000 square feet (112 to 259 pounds/acre) in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 5.9 pounds / 1000 square feet ([462] 518 pounds/acre) per year.


APPLICATION TIMING AND INFORMATION: This product may be applied to newly transplanted and established ornamentals as a broadcast, over the top or directed application. Around new transplants, delay application to allow soil to settle and water thoroughly before applying. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when applied to soil free of clods, weeds, and debris such as leaves. Prior to application, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

TOLERANT ORNAMENTAL SPECIES: This product will not harm most trees, shrubs, vines and flowers. The species listed below* are tolerant to this product. Best results will be obtained when product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.
 (*List may be truncated to save space)

SCIENTIFIC NAME

COMMON NAME

- Abies spp Fir species** (Balsam, Frasier, Noble, etc.)
- Abelia grandiflora Abelia: Sherwood
- Acer palmatum Japanese Maple
- A. platanoides Norway Maple
- Achillea spp. Yarrow, King Edward
- Actinidia chinensis Kiwi*
- Agapanthus africanus Lily-of-the-Nile (African Lily)
- Agapanthus orientalis
- Akebla quinata Five-Leaf or Chocolate Vine
- Allium cemuum Lady's Leek, Nodding Onion
- Anemone hydrida Japanese Anemone
- Aquilegia spp. Aquilegia: Red and Gold
- Arctostaphylos densiflora Vine Hill Manzanita
- Arctotheca calendula Cape Weed
- Artemisia spp. Wormwood: Silver Mound, Castle
- Aster spp. Aster: Bonnie Blue, Purple Dome
- Aster X Frikartii
- Athrium Fitiz-femina Lady Fern; Fern Lady
- Aucuba japonica Japanese Aucuba
- Begonia spp. Fibrous Begonia: Hardy Grandis
- Berberis gladwynensis Barberry
- B. Julianae Wintergreen Barberry
- B. mentorensis Mentor Barberry
- B. thunbergii Japanese Barberry
- B. verruculosa Warty Barberry



14/18

<i>I. opaca</i>	American Holly
<i>I. vomitoria</i>	Yaupon Holly, Schillings
<i>Inula ensifolia</i>	
<i>Iris ensata</i>	Sword-Leaved Iris; Jodlesong
<i>Iris siberica</i>	Siberian Iris; Cabernet
<i>Iris spp.</i>	Iris species**
<i>Jasminium nudiflorum</i>	Winter Jasmine
<i>Juniperus chinensis</i>	Chinese Juniper; Nick's Compact; Parsonii
<i>J. conferta</i>	Shore Juniper; Blue Pacific
<i>J. davuria</i>	Parsoni
<i>J. horizontalis</i>	Creeping Juniper
<i>Juglans spp.</i>	Walnut*
<i>Justicia brandegeana</i>	Shrimp Plant
<i>Lagerstromia indica</i>	Crape Myrtle
<i>Lagerstromia indica X Fauriei</i>	Crape Myrtle, Tuscarora
<i>Lantana montevidensis</i>	Weeping Lantana
<i>Lavender spp.</i>	Lavender; Munstead
<i>Leonopodium alpinum</i>	Edelweiss
<i>Ligustrum amurense</i>	Amur Privet
<i>Ligustrum sinense</i>	Chinese privet, Variiegata
<i>L. japonicum</i>	Japanese Privet
<i>L. lucidum</i>	Glossy Privet (Waxleaf)
<i>Lillium spp.</i>	Lily; Jazz
<u>SCIENTIFIC NAME</u>	<u>COMMON NAME</u>
<i>Liriope muscari</i>	Big Blue Lily; Silver Mound, Evergreen Giant
<i>Liriope muscari var. variegata</i>	Liriope, variegated
<i>Liriope spicata</i>	Liriope, Creeping
<i>Lobelia Cardinalis</i>	Cardinal Flower, Indian Pink
<i>Lonicera japonica</i>	Japanese Honeysuckle
<i>L. tatarica</i>	Tatarian Honeysuckle
<i>Lorapetalum chinense</i>	Burgundy
<i>Lythrum spp.</i>	Loosestrife; Modern Pink
<i>Magnolia spp.</i>	Magnolia species**
<i>Maleophora luteola</i>	Ice Plant
<i>Malus sp.</i>	Crabapple*
<i>Miscanthus sinensis</i>	Yaku Jima, Silberfeder
<i>Nandina domestica</i>	Heavenly Bamboo
<i>Narcissus spp.</i>	Narcissus species**
<i>Nerium oleander</i>	Oleander
<i>Olea europaea</i>	Olive*
<i>Ophiopogon japonicus</i>	Mondo Grass
<i>Osmanthus heterophyllus</i>	Osmanthus (False Holly): Gulf Tide
<i>Osteospermum fruticosum</i>	Trailing African Daisy
<i>Oxydendrum arboreum</i>	Sourwood
<i>Paeonia suffruticosa</i>	Tree Peony
<i>Pennisetumsetaceumi</i>	Fountain Grass (Dwarf)
<i>Perovskia atriplicifolia</i>	
<i>Persea americana</i>	Avocado*
<i>Photinia fraseri</i>	Frasier's Photinia (Redtip)
<i>Physotegia Virginiana</i>	Dragonhead, False; Vivid
<i>Picea spp.</i>	Spruce species** (Colorado Blue, Norway)
<i>Pieris japonica</i>	Lily-of-the-Valley Bush
<i>Pinus brutia</i>	Calabrian Pine
<i>P. canariensis</i>	Canary Island Pine
<i>P. elliottii</i>	Slash Pine
<i>P. halepensis</i>	Aleppo Pine
<i>P. nigra</i>	Austrian Pine
<i>P. radiata</i>	Monterey Pine
<i>P. strobus</i>	Eastern White Pine
<i>P. sylvestris</i>	Scotch Pine

16/18

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal. **PESTICIDE STORAGE:** Store in original container in a locked storage area out of reach of children and domestic animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Nonrefillable container: Do not reuse or refill this container. Offer for recycling, if available. If not available then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Warranty

The directions for use of this product are believed to be adequate and must be followed carefully, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result due to such factors as weather conditions, presence or absence of other materials, or the manner of use or application, all of which are beyond the control of Howard Johnson's Enterprises, Inc. the manufacturer, or the seller. To the extent consistent with applicable law, the products sold to you are furnished "as is" by Howard Johnson's Enterprises, Inc. The manufacturer and the seller are subject only to the manufacturer's warranties, if any, which appear on the label of the product sold to you. Except as warranted by this label Howard Johnson's Enterprises, Inc, the manufacturer, or the seller makes no warranties, guarantees, or representations of any kind to the buyer or the user, either express or implied, or by usage of trade, statutory or otherwise, with regard to the product sold or use of the product, including, but not limited to, merchantability, fitness for a particular purpose or use, or eligibility of the product for any particular trade usage. To the extent consistent with applicable law, Buyer's or user's exclusive remedy, and Howard Johnson's Enterprises, Inc the manufacturer's or the seller's total liability shall be limited to damages not exceeding the cost of the product. No agent or employee of Howard Johnson's Enterprises, Inc, or the seller is authorized to amend the terms of this warranty disclaimer or the product's label or to make a presentation or recommendation different from or inconsistent with the label of this product.

To the extent consistent with applicable law, Howard Johnson's Enterprises, Inc, the manufacturer, or the seller shall not be liable for consequential, special, or indirect damages resulting from the use, handling, application, storage, or disposal of this product or for damages in the nature of penalties, and the buyer and the user waive any right that they may have to such damages.

OPTIONAL STATEMENTS

(Optional wording required by some states in order to effect fertilizer registration:)

When soil testing or professional advice indicates the need for additional plant nutrients, make additional applications of fertilizer products over the growing season. A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing (No of lbs) pounds of nitrogen per 1,000 square feet and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.htm> [OR by calling [1-877-726-6275]]

When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials. Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.htm>.

Information regarding the raw materials composing this product can be obtained by writing to the Guarantor listed on this label and referring to the batch number found on this container.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk or street, back onto your lawn/garden.

17/18

APPLICATION LIMITS: Certain states have limits on fertilizer application rates and timing. Consult your local State Cooperative Extension Service for current information on application limits for [turf/lawn] fertilizers in your area.

(Optional wording for use with fertilizer inert ingredients that cause staining:)

This product may stain driveways, walks, patios, pools, etc. Wash or sweep off thoroughly immediately after each use.

(Optional statement for Proposition 65 in California)

NOTICE- This product contains chemicals known to the State of California to cause cancer and birth defects or other reproductive harm."

(Optional wording in Florida for non-specialty packaging)

In the state of Florida, this product should be applied at rates not exceeding those recommended in the document titled Best Management Practices for Protection of Water Resources in Florida, June 2002, Florida Green Industries.

(Note: If registered in the state of Florida, any of the following optional statements may be added as appropriate) In Florida, do not exceed X applications per year on Bahiagrass in the North, and X per year in Central and South Florida.

In Florida, do not exceed X applications per year on Bermudagrass in the North, or X applications per year in Central and 7 applications per year in South Florida.

In Florida, do not exceed X applications per year on Centipedegrass in the North, or X applications per year in Central and South Florida.

In Florida, do not exceed X applications per year on St. Augustinegrass in the North, or X applications per year in Central and 6 applications per year in South Florida.

In Florida, do not exceed X applications per year on Zoysiagrass in the North, or X applications per year in Central and South Florida.

(Note: may also be expressed in tabular format.)

If Registered in Florida:

Apply only to activity growing turf. Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands in the driveway, sidewalk, or street, back onto your lawn. Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices. [Check with your county or city government to determine if there are local regulations for fertilizer use. / Check with your county or city government to determine if there are local restrictions on fertilizer use or application.]

(Optional Marketing Claims)

Herbicide Benefit Claims

STOPS MANY WEED GRASSES BEFORE THEY GROW

3-5 month Weed Grass Control

Up to 5 [month] [months of] weed grass control

Controls [prevents] crabgrass [and other listed weed grasses] for up to 5 months

Covers [5,000 / 7,500 / 10,000 / 15,000] Sq. Ft.

Prevents crabgrass and other weed grasses

Controls Crabgrass and Broadleaf Weeds Before they Germinate

Prevent crabgrass and fertilize your lawn in one application

Controls Annual Bluegrass (poa annua), Chickweed, Crabgrass, Spurge [and many other weeds]

cccc
cccc
cc c
c c c
c c c
cccccc
c c c
cccccc
c c c
cccccc

Fertilizer Benefit Claims

Enhanced Efficiency Fertilizer [- feed more lawn with less fertilizer]

Delivers nutrients more efficiently than ordinary fertilizers

Quick Greening and Extended Feeding

Helps strengthen lawn against heat & drought

Good for all grass types

[Feeds] [greens] [feeds and greens] for [up to] [3] [X] months

c c c
c c c c
c c c c
c c c c
c c c c
c c c c
c c c c

Contains slow release Nitrogen for extended [greening, feeding]
 [Rich in] Nitrogen for a lush green lawn
 [Enriched with] Potassium for overall plant vigor
 Contains potassium for drought & disease resistance
 Phosphorus Free – [Helping Reduce Phosphorus in Water Sources]
 Provides Lush, green grass
 Promotes thick, full grass
 You only need [XX] pounds of fertilizer to cover (XXXX) square feet

Fungi Benefit Claims

Boosts Nutrient Uptake for a Lush, Green Lawn
 Increases Drought Tolerance & Reduces Watering Needs by Extending Root Absorption Area
 Increased root absorption area for more efficient use of water in [the] soil
 Allows grass to use more of the fertilizer you buy and apply
 [Contains mycorrhizal fungi for] enhanced nutrient uptake
 Increased root absorption area to protect against heat & drought
 Increase absorption [area of roots, of fertilizer]
 The only brand with MYCO Advantage™ for better results than [fertilizer alone, ordinary fertilizer]
 MYCO Advantage™ - Better Results than fertilizer alone

Additional Copy

The Pennington® UltraGreen® Difference

Pennington® UltraGreen® products are more efficient than other fertilizers. Half of the nitrogen in this premium line is stabilized**, which allows grass to use more of the nutrients you put down and reduces nutrient loss to the environment. The other half of the nitrogen is slow-release for extended feeding for [up to] 3 months.

Additionally, these unique blends contain mycorrhizal fungi, natural organisms that have been present in vigorous soils for millions of years. These microscopic fungi increase the surface absorbing area of roots, greatly improving the ability of the lawn to access vital soil resources such as water and nutrients.

These tiny fertilizer granules are packed with essential plant nutrients and fungi, so you can feed more lawn with less fertilizer.

Benefits of Pennington® UltraGreen™ Crabgrass Preventer Plus Fertilizer 30-0-4

- Promotes thick, green grass
- Half of the total nitrogen in this product is slow-release and feeds for [up to] 3 months [guaranteed]
- Half of the total nitrogen is stabilized for less leaching into the environment
- Phosphorus-free

Pennington® UltraGreen® Crabgrass Preventer Plus Fertilizer 30-0-4 WILL NOT BURN when used as directed. We're so confident you'll love the results of using [this product] that it comes with the Pennington Satisfaction Guarantee. †

† Pennington Satisfaction Guarantee: If, after following the instructions, you are not satisfied with the product results, send your name, address, and proof of purchase (original dated receipt) to:

Pennington Refund
 1000 Parkwood Circle, Suite 700
 Atlanta, GA 30339